


FIGURE 4–1a. STEPWISE APPROACH FOR MANAGING ASTHMA IN CHILDREN 0–4 YEARS OF AGE



Key: **Alphabetical order is used when more than one treatment option is listed within either preferred or alternative therapy.** ICS, inhaled corticosteroid; LABA, inhaled long-acting beta₂-agonist; SABA, inhaled short-acting beta₂-agonist

FIGURE 4–3a. ASSESSING ASTHMA CONTROL AND ADJUSTING THERAPY IN CHILDREN 0–4 YEARS OF AGE

Components of Control		Classification of Asthma Control (0–4 years of age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week	>2 days/week	Throughout the day
	Nighttime awakenings	≤1x/month	>1x/month	>1x/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	2–3/year	>3/year
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment (See figure 4–1a for treatment steps.)		<ul style="list-style-type: none">• Maintain current treatment.• Regular followup every 1–6 months.• Consider step down if well controlled for at least 3 months.	<ul style="list-style-type: none">• Step up (1 step) and Reevaluate in 2–6 weeks.• If no clear benefit in 4–6 weeks, consider alternative diagnoses or adjusting therapy.• For side effects, consider alternative treatment options.	<ul style="list-style-type: none">• Consider short course of oral systemic corticosteroids.• Step up (1–2 steps), and Reevaluate in 2 weeks.• If no clear benefit in 4–6 weeks, consider alternative diagnoses or adjusting therapy.• For side effects, consider alternative treatment options.

Key: EIB, exercise-induced bronchospasm


FIGURE 4–2a. CLASSIFYING ASTHMA SEVERITY AND INITIATING TREATMENT IN CHILDREN 0–4 YEARS OF AGE

Assessing severity and initiating therapy in children who are not currently taking long-term control medication

Components of Severity		Classification of Asthma Severity (0–4 years of age)			
		Intermittent	Persistent		
			Mild	Moderate	Severe
Impairment	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	0	1–2x/month	3–4x/month	>1x/week
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2 exacerbations in 6 months requiring oral systemic corticosteroids, or ≥4 wheezing episodes/1 year lasting >1 day AND risk factors for persistent asthma		
		← Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time. →			
		Exacerbations of any severity may occur in patients in any severity category.			
Recommended Step for Initiating Therapy (See figure 4–1a for treatment steps.)		Step 1	Step 2	Step 3 and consider short course of oral systemic corticosteroids	
		In 2–6 weeks, depending on severity, evaluate level of asthma control that is achieved. If no clear benefit is observed in 4–6 weeks, consider adjusting therapy or alternative diagnoses.			

Key: EIB, exercise-induced bronchospasm

FIGURE 4–1b. STEPWISE APPROACH FOR MANAGING ASTHMA IN CHILDREN 5–11 YEARS OF AGE



Key: **Alphabetical order is used when more than one treatment option is listed within either preferred or alternative therapy.** ICS, inhaled corticosteroid; LABA, inhaled long-acting beta₂-agonist; LTRA, leukotriene receptor antagonist; SABA, inhaled short-acting beta₂-agonist


FIGURE 4–2b. CLASSIFYING ASTHMA SEVERITY AND INITIATING TREATMENT IN CHILDREN 5–11 YEARS OF AGE

Assessing severity and initiating therapy in children who are not currently taking long-term control medication

Components of Severity		Classification of Asthma Severity (5–11 years of age)			
		Intermittent	Persistent		
Impairment	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	≤2x/month	3–4x/month	>1x/week but not nightly	Often 7x/week
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
	Lung function	<ul style="list-style-type: none">• Normal FEV₁ between exacerbations• FEV₁ >80% predicted• FEV₁/FVC >85%	<ul style="list-style-type: none">• FEV₁ = >80% predicted• FEV₁/FVC >80%	<ul style="list-style-type: none">• FEV₁ = 60–80% predicted• FEV₁/FVC = 75–80%	<ul style="list-style-type: none">• FEV₁ <60% predicted• FEV₁/FVC <75%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year (see note)	≥2/year (see note)		
		Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time for patients in any severity category. Relative annual risk of exacerbations may be related to FEV ₁ .			
Recommended Step for Initiating Therapy (See figure 4–1b for treatment steps.)		Step 1	Step 2	Step 3, medium-dose ICS option	Step 3, medium-dose ICS option, or step 4
		and consider short course of oral systemic corticosteroids			
In 2–6 weeks, evaluate level of asthma control that is achieved, and adjust therapy accordingly.					

Key: EIB, exercise-induced bronchospasm; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; ICS, inhaled corticosteroids

FIGURE 4–5. STEPWISE APPROACH FOR MANAGING ASTHMA IN YOUTHS ≥12 YEARS OF AGE AND ADULTS



Key: Alphabetical order is used when more than one treatment option is listed within either preferred or alternative therapy. EIB, exercise-induced bronchospasm; ICS, inhaled corticosteroid; LABA, long-acting inhaled beta₂-agonist; LTRA, leukotriene receptor antagonist; SABA, inhaled short-acting beta₂-agonist

FIGURE 4–3b. ASSESSING ASTHMA CONTROL AND ADJUSTING THERAPY IN CHILDREN 5–11 YEARS OF AGE

Components of Control		Classification of Asthma Control (5–11 years of age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week but not more than once on each day	>2 days/week or multiple times on ≤2 days/week	Throughout the day
	Nighttime awakenings	≤1x/month	≥2x/month	≥2x/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
	Lung function <ul style="list-style-type: none">• FEV₁ or peak flow• FEV₁/FVC	>80% predicted/ personal best >80%	60–80% predicted/ personal best 75–80%	<60% predicted/ personal best <75%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year (see note)	
		Consider severity and interval since last exacerbation		
	Reduction in lung growth	Evaluation requires long-term followup.		
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment (See figure 4–1b for treatment steps.)		<ul style="list-style-type: none">• Maintain current step.• Regular followup every 1–6 months.• Consider step down if well controlled for at least 3 months.	<ul style="list-style-type: none">• Step up at least 1 step and Reevaluate in 2–6 weeks.• For side effects: consider alternative treatment options.	<ul style="list-style-type: none">• Consider short course of oral systemic corticosteroids,• Step up 1–2 steps, and Reevaluate in 2 weeks.• For side effects, consider alternative treatment options.

Key: EIB, exercise-induced bronchospasm; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity

FIGURE 4–6. CLASSIFYING ASTHMA SEVERITY AND INITIATING TREATMENT IN YOUTHS ≥12 YEARS OF AGE AND ADULTS

Assessing severity and initiating treatment for patients who are not currently taking long-term control medications

Components of Severity		Classification of Asthma Severity ≥12 years of age			
		Intermittent	Persistent		
Impairment	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	≤2x/month	3–4x/month	>1x/week but not nightly	Often 7x/week
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily, and not more than 1x on any day	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
	Lung function	<ul style="list-style-type: none">• Normal FEV₁ between exacerbations• FEV₁ >80% predicted• FEV₁/FVC normal	<ul style="list-style-type: none">• FEV₁ >80% predicted• FEV₁/FVC normal	<ul style="list-style-type: none">• FEV₁ >60% but <80% predicted• FEV₁/FVC reduced 5%	<ul style="list-style-type: none">• FEV₁ <60% predicted• FEV₁/FVC reduced >5%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year (see note)	≥2/year (see note)		
		Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time for patients in any severity category. Relative annual risk of exacerbations may be related to FEV ₁ .			
Recommended Step for Initiating Treatment (See figure 4–5 for treatment steps.)		Step 1	Step 2	Step 3	Step 4 or 5
		and consider short course of oral systemic corticosteroids			
		In 2–6 weeks, evaluate level of asthma control that is achieved and adjust therapy accordingly.			

Key: FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; ICU, intensive care unit

FIGURE 4–7. ASSESSING ASTHMA CONTROL AND ADJUSTING THERAPY IN YOUTHS ≥12 YEARS OF AGE AND ADULTS

Components of Control		Classification of Asthma Control (≥12 years of age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week	>2 days/week	Throughout the day
	Nighttime awakenings	≤2x/month	1–3x/week	≥4x/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
	FEV ₁ or peak flow	>80% predicted/ personal best	60–80% predicted/ personal best	<60% predicted/ personal best
	Validated questionnaires ATAQ ACQ ACT	0 ≤0.75* ≥20	1–2 ≥1.5 16–19	3–4 N/A ≤15
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year (see note)	
		Consider severity and interval since last exacerbation		
	Progressive loss of lung function	Evaluation requires long-term followup care		
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment (see figure 4–5 for treatment steps)		<ul style="list-style-type: none">• Maintain current step.• Regular followups every 1–6 months to maintain control.• Consider step down if well controlled for at least 3 months.	<ul style="list-style-type: none">• Step up 1 step and Reevaluate in 2–6 weeks.• For side effects, consider alternative treatment options.	<ul style="list-style-type: none">• Consider short course of oral systemic corticosteroids.• Step up 1, 2 steps, and Reevaluate in 2 weeks.• For side effects, consider alternative treatment options.

■
— *ACQ values of 0.76–1.4 are indeterminate regarding well-controlled asthma.
— Key: EIB, exercise-induced bronchospasm; ICU, intensive care unit