

Measure	Patient Population	Compliance Description	Codes
Asthma Medication Ratio	<p>Patients having <u>persistent</u> asthma in both the <u>measurement year</u> and <u>prior year</u>, defined with one of the following:</p> <ul style="list-style-type: none"> At least 1 ED visit with a primary diagnosis of asthma At least 1 acute inpatient discharge with a primary diagnosis of asthma At least 1 acute inpatient discharge with a primary diagnosis of asthma on the discharge claim At least 4 outpatient visits /observation visits *(all 4 any type) with a diagnosis of asthma (all diagnoses) and at least two asthma medication dispensing events for any controller or reliever medication At least four asthma medication dispensing events for any controller or reliever medication <p>If only leukotriene modifiers or antibody inhibitors dispensed, must also have at least 1 asthma diagnosis</p>	<p>The percentage of patients age 5-18 years who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications ≥ 0.50 during the measurement year.</p> <p>Exclusions: Patients with Acute Respiratory Failure, COPD, Conditions related to Fumes/Vapors, Cystic Fibrosis, Emphysema, Obstructive Chronic Bronchitis</p>	<p>Diagnosis: Persistent: J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52 Intermittent: J45.20, J45.21, J45.22 Other: J45.9, J45.901, J45.902, J45.909, J45.99, J45.990, J45.991, J45.998</p>
Chlamydia Screening in Women	<p>Women age 16-24 years during the measurement year who are sexually active. Sexual activity is identified using 2 methods: pharmacy and claims/encounter data. (The organization must use both methods to identify the eligible population but only needs to be identified in one method to be eligible for the measure.)</p> <p>Claim encounter data-indicating sexual activity during measurement year. A code from the following meets criteria:</p> <ol style="list-style-type: none"> 1) Pregnancy Value set 2) Sexual Activity Value set 3) Pregnancy Test Value set <p>Pharmacy data –Patients who were dispensed prescription contraceptives during the measurement year including Diaphragm and spermicide</p>	<p>Evidence of at least one test for chlamydia during the measurement year.</p> <p>Exclusions: Patient excluded if qualified for the denominator based on a pregnancy test alone and meets one of the following:</p> <ol style="list-style-type: none"> 1. A pregnancy test during the measurement year and a prescription for isotretinoin on the date of the pregnancy test or the six days after the pregnancy test. 2. A pregnancy test during the measurement year and an x-ray on the date of the pregnancy test or the six days after the pregnancy test. 	<p>CPT: 87110, 87270, 87320, 87490-87492, 87810</p>
Depression Screening	<p>Patients age 12-18 years</p>	<p>Evidence of a depression screening with any provider during the measurement year</p> <p>Exclusions: 1. Hospice, ever 2. Any diagnosis of depression, bipolar, or delirium before the qualifying depression screen in measurement year or year prior to the measurement year</p>	<p>CPT: 96127</p>
Early & Periodic Screening Participant Ratio (0-6 Years)	<p>Patients age 0-6 years</p>	<p>Evidence of at least 1 EPSDT visit with a primary care provider (pediatrics, family medicine, nurse practitioner, physician assistant) during the measurement year</p>	

Measure	Patient Population	Compliance Description	Codes
Age 2 Immunizations Combo-10	Patients who turn 2 years of age in the measurement year. 12 months of continuous enrollment - no more than 1 gap of up to 45 days during the 12 months prior to the child's 2nd birthday	Evidence of receiving the following by 2 years of age: - 4 DTaP - 3 IPV - 1 MMR - 3 HiB - 3 Hepatitis B - 4 PCV - 1 VZV - 2 or 3 Rotavirus (2 vs. 3 dependent on vaccine series) - 1 Hepatitis A - 2 Influenza Exclusions: Patients with immunodeficiency, HIV, Lymphoreticular cancer, multiple myeloma or leukemia, and anaphylactic reaction to Neomycin.	Diagnosis: Z23 CPT: DTaP – 90723, 90700, 90698, 90696 IPV – 90723, 90698, 90713, 90696 MMR – 90707, 90710 HiB – 90647, 90648, 90698 Hepatitis B – 90723, 90744 PCV – 90670 VZV – 90716, 90710 Rotavirus – 90680, 90681 Hepatitis A – 90633, 90636 Influenza – 90686, 90685, 90672, 90688, 90660, 90656, 90662, 90655, 90658, 90687
Immunizations Childhood Age 6	Patients who turn 6 years of age in the measurement year	Evidence of receiving the following between the 4th and the 6th birthday: 1 DTaP 1 IPV 1 MMR 1 VZV 1 Influenza between 5th & 6th birthday	Diagnosis: Z23 CPT: DTaP – 90723, 90700, 90698, 90696 IPV – 90723, 90698, 90713, 90696 MMR – 90707, 90710 VZV – 90716, 90710 Influenza – 90686, 90685, 90672, 90688, 90660, 90656, 90662, 90655, 90658, 90687
Age 13 Immunizations (MCV, TDAP)	Patients who turn 13 years of age in the measurement year. Continuous enrollment of 12 months prior to the patients 13th birthday.	Evidence of receiving the following: - 1 MCV (between 11th and 13th birthday) - 1 Tdap/Td or 1 Tetanus/1 Diphtheria (between 10th and 13th birthday) Exclusions: Optional exclusion criteria: Anaphylactic reaction to the vaccine or its components any time on or before the child's 13th birthday	Diagnosis: Z23 CPT: MCV – 90734, 90620, 90621, 90733 Tdap/TD/Tetnus/Diphtheria – 90715, 90723, 90700, 90698, 90696, 90714 Influenza – 90686, 90685, 90672, 90688, 90660, 90656, 90662, 90655, 90658, 90687
Age 13 Immunizations (MCV, TDAP, HPV)	Patients who turn 13 years of age in the measurement year. Continuous enrollment of 12 months prior to the patients 13th birthday.	Evidence of receiving the following: - 1 MCV (between 11th and 13th birthday) - 1 Tdap/Td or 1 Tetanus/1 Diphtheria (on or between 10th and 13th birthday) - 2 HPV (between 9th and 13th birthday) (Dates of service must be at least 146 days apart) Exclusions: Optional exclusion criteria: Anaphylactic reaction to the vaccine or its components any time on or before the child's 13th birthday	Diagnosis: Z23 CPT: 90649, 90650, 90651
Lead Screening in Children	Patients who turn 2 years of age during the measurement year	Evidence of at least 1 lead capillary or venous blood test on or before child's second birthday.	CPT: 83655
Social Determinants of Health (SDoH) Screening	Patients age 0-21 years with at least one visit with a primary care provider in the measurement year. No more than one gap in enrollment of up to 45 days during the measurement year	Evidence of at least one SDoH screened in the measurement year. Patients must be screening using standardized SDoH screening questions (at least but not limited to Transportation, Food Insecurity, & Housing).	G9920 - Screening Performed and Negative G9919 - Screening Performed and Positive and Provision of Recommendations

Measure	Patient Population	Compliance Description	Codes
Timeliness of Prenatal Care	Patients pregnant during the measurement year; no specified age	The percentage of deliveries that received a prenatal care visit as a patient of the organization in the first trimester, on the enrollment start date or within 42 days of enrollment in the organization. Exclusions: In Hospice; Non-live births	
Tobacco Use Screening (For more detail see Coding Reference Guide)	Patients age 13-18 years who are non-tobacco users or whose tobacco status is unknown	Evidence of at least one tobacco screening with any provider during the measurement year or year prior	M Codes: 1000F – tobacco use assessed 1034F – tobacco user, smoker 1035F – tobacco user, chew/snuff
Tobacco Cessation Intervention (For more detail see Coding Reference Guide)	Patients age 13-18 years who are current tobacco users in the measurement year or prior year	Evidence of cessation counseling by any provider during the measurement year or year prior	Diagnosis: Z71.6 CPT: 99406 M Code: 4000F
Weight Assessment BMI Percentile (For more detail see Coding Reference Guide)	Patients age 3-17 years. Starts at age 2 years and 1 day.	Evidence of BMI percentile documented Exclusions: Female patients who have a diagnosis of pregnancy during the measurement year	Diagnosis: Z68.51, Z68.52, Z68.53, Z68.54
Weight Assessment Counseling for Nutrition (For more detail see Coding Reference Guide)	Patients age 3-17 years in the measurement year who have had an outpatient visit	Evidence of counseling for nutrition	Diagnosis: Z71.3 CPT: 97802, 97803
Well-Child Visits First 15 Months of Life	Patients who turn 15 months of age during the measurement year. Calculate the 15-month birthday as the child's first birthday plus 90 days.	Evidence of 6 or more well visits with a primary care provider (pediatrics, family medicine, nurse practitioner, physician assistant) within the first 15 months of a child's life	Diagnosis: Z00.110, Z00.111, Z00.121, Z00.129, Z02.89 CPT: 99381, 99382, 99391, 99392
Well-Child Visits - 15-30 Months	Patients who turn 30 months of age during the measurement year. Calculate the 30-month birthday as the child's second birthday plus 180 days.	Evidence of two or more well-child visits between the child's 15-month birthday plus 1 day and the 30-month birthday	Diagnosis: Z00.110, Z00.111, Z00.121, Z00.129, Z02.89 CPT: 99381, 99382, 99391, 99392
Well-Child Ages 3-11 Years	Patients age 3-11 years during the measurement year	Evidence of at least 1 well-child visit with a primary care provider (pediatrics, family medicine, nurse practitioner, physician assistant) during the measurement year	Diagnosis: Z00.121, Z00.129, Z02.89 CPT: 99382, 99383, 99392, 99393
Well-Child Visits - 12-21 Years of Life	Patients age 12-21 years during the measurement year	Evidence of at least 1 well-child visit with a primary care provider (pediatrics, family medicine, nurse practitioner, physician assistant) during the measurement year	Diagnosis: Z00.00, Z00.121, Z00.129, Z02.89 CPT: 99384, 99385, 99394, 99395

Note: Except where noted above all measures require continuous enrollment with no more than one gap in enrollment up to 45 days.

Note: In most cases Telehealth visits can be used to meet measure compliance.